

Literatur

- Anguelova M, Benkelfat C, Turecki G. A systematic review of association studies investigating genes coding for serotonin receptors and the serotonin transporter: II. Suicidal behavior. *Mol Psychiatry* 2003; 8(7): 646–653.
- Arango V, Ernberger P, Marzuk PM, Chen JS, Tierney H, Stanley M et al. Autoradiographic demonstration of increased serotonin 5-HT₂ and beta-adrenergic receptor binding sites in the brain of suicide victims. *Arch Gen Psychiatry* 1990; 47(11): 1038–1047.
- Arango V, Huang YY, Underwood MD, Mann JJ. Genetics of the serotonergic system in suicidal behavior. *J Psychiatr Res* 2003; 37(5): 375–386.
- Arango V, Underwood MD, Mann JJ. Fewer pigmented locus coeruleus neurons in suicide victims: preliminary results. *Biol Psychiatry* 1996; 39(2): 112–120.
- Arango V, Underwood MD, Mann JJ. Postmortem findings in suicide victims. Implications for in vivo imaging studies. *Ann N Y Acad Sci* 1997; 836: 269–287.
- Asberg M. Neurotransmitters and suicidal behavior. The evidence from cerebrospinal fluid studies. *Ann N Y Acad Sci* 1997; 836: 158–181.
- Baldessarini RJ, Hennen J. Genetics of suicide: an overview. *Harv Rev Psychiatry* 2004; 12(1): 1–13.
- Bellivier F, Chaste P, Malafosse A. Association between the TPH gene A218C polymorphism and suicidal behavior: a meta-analysis. *Am J Med Genet B Neuropsychiatr Genet* 2004; 124B(1): 87–91.
- Biegion A, Fieldust S. Reduced tyrosine hydroxylase immunoreactivity in locus coeruleus of suicide victims. *Synapse* 1992; 10(1): 79–82.
- Bondy B, Baghai TC, Minov C, Schule C, Schwarz MJ, Zwanzger P et al. Substance P serum levels are increased in major depression: preliminary results. *Biol Psychiatry* 2003; 53(6): 538–542.
- Bondy B, Buettner A, Zill P. Genetics of suicide. *Mol Psychiatry* 2006; 11(4): 336–351.
- Bondy B, Kuznik J, Baghai T, Schule C, Zwanzger P, Minov C et al. Lack of association of serotonin-2A receptor gene polymorphism (T102C) with suicidal ideation and suicide. *Am J Med Genet* 2000; 96(6): 831–835.
- Brent DA, Johnson BA, Perper J, Connolly J, Bridge J, Bartle S et al. Personality disorder, personality traits, impulsive violence, and completed suicide in adolescents. *J Am Acad Child Adolesc Psychiatry* 1994; 33(8): 1080–1086.
- Brent DA, Mann JJ. Family genetic studies, suicide, and suicidal behavior. *Am J Med Genet C Semin Med Genet* 2005; 133C(1): 13–24.
- Bronisch T, Brunner J, Bondy B, Rujescu D, Bishof G, Heuser I, Müller-Oerlinghausen B, Hawellek B, Maier W, Rao ML, Felber W, Lewitzka U, Oehler J, Broocks A, Hohagen F, Lauterbach E. A multicenter study about Neurobiology of Suicidal Behavior: design, development, and preliminary results. *Arch Suicide Res* 2005; 9(1): 19–26.
- Brunner HG, Nelen MR, van Zandvoort P, Abeling NG, van Gennip AH, Wolters EC et al. X-linked borderline mental retardation with prominent behavioral disturbance: phenotype, genetic localization, and evidence for disturbed monoamine metabolism. *Am J Hum Genet* 1993; 52(6): 1032–1039.
- Brunner J, Bronisch T. [Neurobiological correlates of suicidal behavior]. *Fortschr Neurol Psychiatr* 1999; 67(9): 391–412.
- Brunner J, Parhofer KG, Schwandt P, Bronisch T. Cholesterol, essential fatty acids, and suicide. *Pharmacopsychiatry* 2002; 35(1): 1–5.
- Buchsbaum MS, Coursey RD, Murphy DL. The biochemical high-risk paradigm: behavioral and familial correlates of low platelet monoamine oxidase activity. *Science* 1976; 194(4262): 339–341.
- Burgess DL, Kohrman DC, Galt J, Plummer NW, Jones JM, Spear B et al. Mutation of a new sodium channel gene, *Scn8a*, in the mouse mutant 'motor endplate disease'. *Nat Genet* 1995; 10(4): 461–465.
- Cao L, Li T, Xu K, Liu X. [Association analysis of attempted suicide and 5-HT_{2A} receptor gene]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 2001; 18(3): 195–197.
- Caspi A, McClay J, Moffitt TE, Mill J, Martin J, Craig IW et al. Role of genotype in the cycle of violence in maltreated children. *Science* 2002; 297(5582): 851–854.
- Coursey RD, Buchsbaum MS, Murphy DL. Platelet MAO activity and evoked potentials in the identification of subjects biologically at risk for psychiatric disorders. *Br J Psychiatry* 1979; 134: 372–381.
- Crawford J, Sutherland GR, Goldney RD. No evidence for association of 5-HT_{2A} receptor polymorphism with suicide. *Am J Med Genet* 2000; 96(6): 879–880.
- Crawford J, Zielinski MA, Fisher LJ, Sutherland GR, Goldney RD. Is there a relationship between Wolf-ram syndrome carrier status and suicide? *Am J Med Genet* 2002; 114(3): 343–346.
- De Luca V, Mueller DJ, Tharmalingam S, King N, Kennedy JL. Analysis of the novel TPH2 gene in bipolar disorder and suicidality. *Mol Psychiatry* 2004; 9(10): 896–897.
- De Luca V, Voineskos D, Wong GW, Shinkai T, Rothe C, Strauss J et al. Promoter polymorphism of second tryptophan hydroxylase isoform (TPH2) in schizophrenia and suicidality. *Psychiatry Res* 2005; 134(2): 195–198.
- Dervic K, Grunbaum ME, Burke AK, Mann JJ, Oquendo MA. Cluster C personality disorders in major depressive episodes: the relationship between hostility and suicidal behavior. *Arch Suicide Res* 2007; 11(1): 83–90.
- Dietrich PS, McGivern JG, Delgado SG, Koch BD, Eglon RM, Hunter JC et al. Functional analysis of a voltage-gated sodium channel and its splice variant from rat dorsal root ganglia. *J Neurochem* 1998; 70(6): 2262–2272.
- Du L, Bakish D, Lapierre YD, Ravindran AV, Hrdina PD. Association of polymorphism of serotonin 2A receptor gene with suicidal ideation in major depressive disorder. *Am J Med Genet* 2000; 96(1): 56–60.
- Du L, Faludi G, Palkovits M, Sotonyi P, Bakish D, Hrdina PD. High activity-related allele of MAO-A gene associated with depressed suicide in males. *Neuroreport* 2002; 13(9): 1195–1198.
- Dwivedi Y, Rizavi HS, Conley RR, Roberts RC, Tamminga CA, Pandey GN. Altered gene expression of brain-derived neurotrophic factor and receptor tyrosine kinase B in postmortem brain of suicide subjects. *Arch Gen Psychiatry* 2003; 60(8): 804–815.
- Fenster SD, Chung WJ, Zhai R, Cases-Langhoff C, Voss B, Garner AM et al. Piccolo, a presynaptic zinc finger protein structurally related to bassoon. *Neuron* 2000; 25(1): 203–214.
- Finckh U, Rommelspacher H, Kuhn S, Dufeu P, Otto G, Heinz A et al. Influence of the dopamine D2 receptor (DRD2) genotype on neuroadaptive effects of alcohol and the clinical outcome of alcoholism. *Pharmacogenetics* 1997; 7(4): 271–281.
- Frances A, Fyer M, Clarkin J. Personality and suicide. *Ann N Y Acad Sci* 1986; 487: 281–293.
- Geijer T, Frisch A, Persson ML, Wasserman D, Rockah R, Michaelovsky E et al. Search for association between suicide attempt and serotonergic polymorphisms. *Psychiatr Genet* 2000; 10(1): 19–26.
- Giegling I, Hartmann AM, Moller HJ, Rujescu D. Anger- and aggression-related traits are associated with polymorphisms in the 5-HT-2A gene. *J Affect Disord* 2006; 96(1–2): 75–81.
- Giegling I, Moreno-De-Luca D, Rujescu D, Schneider B, Hartmann AM, Schnabel A et al. Dopa decarboxylase and tyrosine hydroxylase gene variants in suicidal behavior. *Am J Med Genet B Neuropsychiatr Genet* 2008; 147(3): 308–315.
- Giegling I, Rujescu D, Mandelli L, Schneider B, Hartmann AM, Schnabel A et al. Tachykinin receptor 1 variants associated with aggression in suicidal behavior. *Am J Med Genet B Neuropsychiatr Genet* 2007; 144B(6): 757–761.
- Giegling I, Rujescu D, Mandelli L, Schneider B, Hartmann AM, Schnabel A et al. Estrogen receptor gene 1 variants are not associated with suicidal behavior. *Psychiatry Res* 2008; 160(1): 1–7.
- Gietl A, Giegling I, Hartmann AM, Schneider B, Schnabel A, Maurer K et al. ABCG1 gene variants in suicidal behavior and aggression-related traits. *Eur Neuropsychopharmacol* 2007; 17(6–7): 410–416.
- Golomb BA. Cholesterol and violence: is there a connection? *Ann Intern Med* 1998; 128(6): 478–487.
- Gross-Isseroff R, Biegion A, Voet H, Weizman A. The suicide brain: a review of postmortem receptor/transporter binding studies. *Neurosci Biobehav Rev* 1998; 22(5): 653–661.
- Gwadry FG, Sequeira A, Hoke G, Ffrench-Mullen JM, Turecki G. Molecular characterization of suicide by microarray analysis. *Am J Med Genet C Semin Med Genet* 2005; 133C(1): 48–56.
- Hallman J, von Knorring L, Orelund L. Personality disorders according to DSM-III-R and thrombocyte monoamine oxidase activity in type 1 and type 2 alcoholics. *J Stud Alcohol* 1996; 57(2): 155–161.
- Hesselbrock V, Dick D, Hesselbrock M, Foroud T, Schuckit M, Edenberg H et al. The search for genetic risk factors associated with suicidal behavior. *Alcohol Clin Exp Res* 2004; 28(5 Suppl): 70S–76S.
- Ho LW, Furlong RA, Rubinsztein JS, Walsh C, Paykel ES, Rubinsztein DC. Genetic associations with clinical characteristics in bipolar affective disorder and recurrent unipolar depressive disorder. *Am J Med Genet* 2000; 96(1): 36–42.
- Hong CJ, Huo SJ, Yen FC, Tung CL, Pan GM, Tsai SJ. Association study of a brain-derived neurotrophic-factor genetic polymorphism and mood disorders, age of onset and suicidal behavior. *Neuropsychobiology* 2003; 48(4): 186–189.
- Hong CJ, Pan GM, Tsai SJ. Association study of onset age, attempted suicide, aggressive behavior, and schizophrenia with a serotonin 1B receptor (A-161T) genetic polymorphism. *Neuropsychobiology* 2004; 49(1): 1–4.

50. Hong CJ, Wang YC, Tsai SJ. Association study of angiotensin I-converting enzyme polymorphism and symptomatology and antidepressant response in major depressive disorders. *J Neural Transm* 2002; 109(9): 1209–1214.
51. Hrdina PD, Demeter E, Vu TB, Sotonyi P, Palkovits M. 5-HT uptake sites and 5-HT₂ receptors in brain of antidepressant-free suicide victims/depressives: increase in 5-HT₂ sites in cortex and amygdala. *Brain Res* 1993; 614(1–2): 37–44.
52. Huang YY, Battistuzzi C, Oquendo MA, Harkavy-Friedman J, Greenhill L, Zalsman G et al. Human 5-HT_{1A} receptor C(-1019)G polymorphism and psychopathology. *Int J Neuropsychopharmacol* 2004; 7(4): 441–451.
53. Huang YY, Cate SP, Battistuzzi C, Oquendo MA, Brent D, Mann JJ. An association between a functional polymorphism in the monoamine oxidase a gene promoter, impulsive traits and early abuse experiences. *Neuropsychopharmacology* 2004; 29(8): 1498–1505.
54. Huang YY, Grailhe R, Arango V, Hen R, Mann JJ. Relationship of psychopathology to the human serotonin 1B genotype and receptor binding kinetics in postmortem brain tissue. *Neuropsychopharmacology* 1999; 21(2): 238–246.
55. Huang YY, Oquendo MA, Friedman JM, Greenhill LL, Brodsky B, Malone KM et al. Substance abuse disorder and major depression are associated with the human 5-HT_{1B} receptor gene (HTR1B) G861C polymorphism. *Neuropsychopharmacology* 2003; 28(1): 163–169.
56. Johann M, Putzhammer A, Eichhammer P, Wodarz N. Association of the -141C Del variant of the dopamine D₂ receptor (DRD2) with positive family history and suicidality in German alcoholics. *Am J Med Genet B Neuropsychiatr Genet* 2005; 132B(1): 46–49.
57. Johns CA, Stanley M, Stanley B. Suicide in schizophrenia. *Ann N Y Acad Sci* 1986; 487: 294–300.
58. Jones G, Zammit S, Norton N, Hamshere ML, Jones SJ, Millham C et al. Aggressive behaviour in patients with schizophrenia is associated with catechol-O-methyltransferase genotype. *Br J Psychiatry* 2001; 179: 351–355.
59. Kashani JH, Goddard P, Reid JC. Correlates of suicidal ideation in a community sample of children and adolescents. *J Am Acad Child Adolesc Psychiatry* 1989; 28(6): 912–917.
60. Koller G, Bondy B, Preuss UW, Zill P, Soyka M. The C(-1019)G 5-HT_{1A} promoter polymorphism and personality traits: no evidence for significant association in alcoholic patients. *Behav Brain Funct* 2006; 2: 7.
61. Koller G, Engel RR, Preuss UW, Karakesisoglou A, Zill P, Bondy B et al. Tryptophan hydroxylase gene 1 polymorphisms are not associated with suicide attempts in alcohol-dependent individuals. *Addict Biol* 2005; 10(3): 269–273.
62. Kotler M, Barak P, Cohen H, Averbuch IE, Grinshpoon A, Gritsenko I et al. Homicidal behavior in schizophrenia associated with a genetic polymorphism determining low catechol O-methyltransferase (COMT) activity. *Am J Med Genet* 1999; 88(6): 628–633.
63. Kramer MS, Winokur A, Kelsey J, Preskorn SH, Rothschild AJ, Snavely D et al. Demonstration of the efficacy and safety of a novel substance P (NK1) receptor antagonist in major depression. *Neuropsychopharmacology* 2004; 29(2): 385–392.
64. Kunugi H, Hashimoto R, Yoshida M, Tatsumi M, Kamijima K. A missense polymorphism (S205L) of the low-affinity neurotrophin receptor p75NTR gene is associated with depressive disorder and attempted suicide. *Am J Med Genet B Neuropsychiatr Genet* 2004; 129B(1): 44–46.
65. Kunugi H, Ishida S, Kato T, Tatsumi M, Sakai T, Hattori M et al. A functional polymorphism in the promoter region of monoamine oxidase-A gene and mood disorders. *Mol Psychiatry* 1999; 4(4): 393–395.
66. Lachman HM, Papolos DF, Saito T, Yu YM, Szumlanski CL, Weinshilboum RM. Human catechol-O-methyltransferase pharmacogenetics: description of a functional polymorphism and its potential application to neuropsychiatric disorders. *Pharmacogenetics* 1996; 6(3): 243–250.
67. Lalovic A, Sequeira A, DeGuzman R, Chawky N, Lesage A, Seguin M et al. Investigation of completed suicide and genes involved in cholesterol metabolism. *J Affect Disord* 2004; 79(1–3): 25–32.
68. Lalovic A, Turecki G. Meta-analysis of the association between tryptophan hydroxylase and suicidal behavior. *Am J Med Genet* 2002; 114(5): 533–540.
69. Lauterbach E, Brunner J, Hawellek B, Lewitzka U, Ising M, Bondy B, Rao ML, Frahnert C, Rujescu D, Müller-Oerlinghausen B, Schley J, Heuser I, Maier W, Hohagen F, Felber W, Bronisch T. Platelet 5-HT_{2A} receptor binding and tryptophan availability in depression are not associated with recent history of suicide attempts but with personality traits characteristic for suicidal behavior. *J Affect Disord* 2006 Mar; 91(1): 57–62.
70. Lemonde S, Turecki G, Bakish D, Du L, Hrdina PD, Bown CD et al. Impaired repression at a 5-hydroxytryptamine 1A receptor gene polymorphism associated with major depression and suicide. *J Neurosci* 2003; 23(25): 8788–8799.
71. Lesch KP, Bengel D, Heils A, Sabol SZ, Greenberg BD, Petri S et al. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science* 1996; 274(5292): 1527–1531.
72. Levitt M, Spector S, Sjoerdsma A, Udenfriend S. Elucidation of the rate-limiting step in norepinephrine biosynthesis in the perfused guinea-pig heart. *J Pharmacol Exp Ther* 1965; 148: 1–8.
73. Lewitzka U, Müller-Oerlinghausen B, Felber W, Brunner J, Hawellek B, Rujescu D et al. Is MAO-B activity in platelets associated with the occurrence of suicidality and behavioural personality traits in depressed patients? *Acta Psychiatr Scand* 2008; 117(1): 41–49.
74. Lin PY, Tsai G. Association between serotonin transporter gene promoter polymorphism and suicide: results of a meta-analysis. *Biol Psychiatry* 2004; 55(10): 1023–1030.
75. Liou YJ, Tsai SJ, Hong CJ, Wang YC, Lai IC. Association analysis of a functional catechol-o-methyltransferase gene polymorphism in schizophrenic patients in Taiwan. *Neuropsychobiology* 2001; 43(1): 11–14.
76. Maier W. Genetik suizidalen Verhaltens. In: Wolfersdorf M, Kaschka WP, editors. Suizidalität. Die biologische Dimension. Berlin, Heidelberg: Springer 1996: 85–95.
77. Mann JJ. The neurobiology of suicide. *Nat Med* 1998; 4(1): 25–30.
78. Mann JJ. A current perspective of suicide and attempted suicide. *Ann Intern Med* 2002; 136(4): 302–311.
79. Mann JJ. Neurobiology of suicidal behaviour. *Nat Rev Neurosci* 2003; 4(10): 819–828.
80. Mann JJ, Apter A, Bertolote J, Beautrais A, Currier D, Haas A et al. Suicide prevention strategies: a systematic review. *JAMA* 2005; 294(16): 2064–2074.
81. Mann JJ, Henteleff RA, Lagattuta TF, Perper JA, Li S, Arango V. Lower 3H-paroxetine binding in cerebral cortex of suicide victims is partly due to fewer high affinity, non-transporter sites. *J Neural Transm* 1996; 103(11): 1337–1350.
82. Mann JJ, Stanley M, McBride PA, McEwen BS. Increased serotonin₂ and beta-adrenergic receptor binding in the frontal cortices of suicide victims. *Arch Gen Psychiatry* 1986; 43(10): 954–959.
83. McGuffin P, Marusic A, Farmer A. What can psychiatric genetics offer suicidology? *Crisis* 2001; 22(2): 61–65.
84. Middendorfer-Rutz E, Rasmussen F, Wasserman D. Familial clustering of suicidal behaviour and psychopathology in young suicide attempters. A register-based nested case control study. *Soc Psychiatry Psychiatr Epidemiol* 2008; 43(1): 28–36.
85. Müller DJ, Barkow K, Kovalenko S, Ohlraun S, Fangerau H, Kölsch H, Lemke MR, Held T, Nöthen MM, Maier W, Heun R, Rietschel M. Suicide attempts in schizophrenia and affective disorders with relation to some specific demographical and clinical characteristics. *Eur Psychiatry* 2005 Jan; 20(1): 65–69.
86. New AS, Gelernter J, Goodman M, Mitropoulou V, Koenigsberg H, Silverman J et al. Suicide, impulsive aggression, and HTR1B genotype. *Biol Psychiatry* 2001; 50(1): 62–65.
87. Nishiguchi N, Shirakawa O, Ono H, Nishimura A, Nushida H, Ueno Y et al. No evidence of an association between 5HT_{1B} receptor gene polymorphism and suicide victims in a Japanese population. *Am J Med Genet* 2001; 105(4): 343–345.
88. Nishiguchi N, Shirakawa O, Ono H, Nishimura A, Nushida H, Ueno Y et al. Lack of an association between 5-HT_{1A} receptor gene structural polymorphisms and suicide victims. *Am J Med Genet* 2002; 114(4): 423–425.
89. Nolan KA, Volavka J, Czobor P, Cseh A, Lachman H, Saito T et al. Suicidal behavior in patients with schizophrenia is related to COMT polymorphism. *Psychiatr Genet* 2000; 10(3): 117–124.
90. Ohara K, Nagai M, Tsukamoto T, Tani K, Suzuki Y, Ohara K. 5-HT_{2A} receptor gene promoter polymorphism--1438G/A and mood disorders. *Neuroreport* 1998; 9(6): 1139–1141.
91. Ohtani M, Shindo S, Yoshioka N. Polymorphisms of the tryptophan hydroxylase gene and serotonin 1A receptor gene in suicide victims among Japanese. *Tohoku J Exp Med* 2004; 202(2): 123–133.
92. Ono H, Shirakawa O, Nishiguchi N, Nishimura A, Nushida H, Ueno Y et al. Serotonin 2A receptor gene polymorphism is not associated with completed suicide. *J Psychiatr Res* 2001; 35(3): 173–176.
93. Ono H, Shirakawa O, Nishiguchi N, Nishimura A, Nushida H, Ueno Y et al. No evidence of an association between a functional monoamine oxidase a gene polymorphism and completed suicides. *Am J Med Genet* 2002; 114(3): 340–342.
94. Ono H, Shirakawa O, Nushida H, Ueno Y, Maeda K. Association between catechol-O-methyltransferase

- functional polymorphism and male suicide completers. *Neuropsychopharmacology* 2004; 29(7): 1374–1377.
95. Ordway GA, Smith KS, Haycock JW. Elevated tyrosine hydroxylase in the locus coeruleus of suicide victims. *J Neurochem* 1994; 62(2): 680–685.
96. Persson ML, Geijer T, Wasserman D, Rockah R, Frisch A, Michaelovsky E et al. Lack of association between suicide attempt and a polymorphism at the dopamine receptor D4 locus. *Psychiatr Genet* 1999; 9(2): 97–100.
97. Persson ML, Wasserman D, Geijer T, Jonsson EG, Terenius L. Tyrosine hydroxylase allelic distribution in suicide attempters. *Psychiatry Res* 1997; 72(2): 73–80.
98. Pitchot W, Hansenne M, Ansseau M. Role of dopamine in non-depressed patients with a history of suicide attempts. *Eur Psychiatry* 2001; 16(7): 424–427.
99. Pitchot W, Hansenne M, Moreno AG, Ansseau M. Suicidal behavior and growth hormone response to apomorphine test. *Biol Psychiatry* 1992; 31(12): 1213–1219.
100. Placidi GP, Oquendo MA, Malone KM, Huang YY, Ellis SP, Mann JJ. Aggressivity, suicide attempts, and depression: relationship to cerebrospinal fluid monoamine metabolite levels. *Biol Psychiatry* 2001; 50(10): 783–791.
101. Pooley EC, Houston K, Hawton K, Harrison PJ. Deliberate self-harm is associated with allelic variation in the tryptophan hydroxylase gene (TPH A779C), but not with polymorphisms in five other serotonergic genes. *Psychol Med* 2003; 33(5): 775–783.
102. Preuss UW, Koller G, Bahlmann M, Soyka M, Bondy B. No association between suicidal behavior and 5-HT_{2A}-T102C polymorphism in alcohol dependents. *Am J Med Genet* 2000; 96(6): 877–878.
103. Pucadyil TJ, Chattopadhyay A. Cholesterol modulates ligand binding and G-protein coupling to serotonin(1A) receptors from bovine hippocampus. *Biochim Biophys Acta* 2004; 1663(1–2): 188–200.
104. Qin P, Agerbo E, Mortensen PB. Suicide risk in relation to family history of completed suicide and psychiatric disorders: a nested case-control study based on longitudinal registers. *Lancet* 2002; 360(9340): 1126–1130.
105. Reif A, Lesch KP. Toward a molecular architecture of personality. *Behav Brain Res* 2003; 139(1–2): 1–20.
106. Reist C, Haier RJ, DeMet E, Chiciz-DeMet A. Platelet MAO activity in personality disorders and normal controls. *Psychiatry Res* 1990; 33(3): 221–227.
107. Rigat B, Hubert C, Alhenc-Gelas F, Cambien F, Corvol P, Soubrier F. An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *J Clin Invest* 1990; 86(4): 1343–1346.
108. Rommelspacher H, May T, Dufeu P, Schmidt LG. Longitudinal observations of monoamine oxidase B in alcoholics: differentiation of marker characteristics. *Alcohol Clin Exp Res* 1994; 18(6): 1322–1329.
109. Roy A. Family history of suicide. *Arch Gen Psychiatry* 1983; 40(9): 971–974.
110. Roy A, Linnoila M. Alcoholism and suicide. *Suicide Life Threat Behav* 1986; 16(2): 244–273.
111. Roy A, Segal NL, Centerwall BS, Robinette CD. Suicide in twins. *Arch Gen Psychiatry* 1991; 48(1): 29–32.
112. Rujescu D, Giegling I, Dahmen N, Szegedi A, Angheliescu I, Gietl A et al. Association study of suicidal behavior and affective disorders with a genetic polymorphism in ABCG1, a positional candidate on chromosome 21q22.3. *Neuropsychobiology* 2000; 42 Suppl 1: 22–25.
113. Rujescu D, Giegling I, Gietl A, Hartmann AM, Moller HJ. A functional single nucleotide polymorphism (V158M) in the COMT gene is associated with aggressive personality traits. *Biol Psychiatry* 2003; 54(1): 34–39.
114. Rujescu D, Giegling I, Mandelli L, Schneider B, Hartmann AM, Schnabel A et al. NOS-I and -III gene variants are differentially associated with facets of suicidal behavior and aggression-related traits. *Am J Med Genet B Neuropsychiatr Genet* 2008; 147B(1): 42–48.
115. Rujescu D, Giegling I, Sato T, Hartmann AM, Moller HJ. Genetic variations in tryptophan hydroxylase in suicidal behavior: analysis and meta-analysis. *Biol Psychiatry* 2003; 54(4): 465–473.
116. Rujescu D, Giegling I, Sato T, Moller HJ. Lack of association between serotonin 5-HT_{1B} receptor gene polymorphism and suicidal behavior. *Am J Med Genet B Neuropsychiatr Genet* 2003; 116B(1): 69–71.
117. Rujescu D, Thalmeier A, Moller HJ, Bronisch T, Giegling I. Molecular genetic findings in suicidal behavior: what is beyond the serotonergic system? *Arch Suicide Res* 2007; 11(1): 17–40.
118. Sabol SZ, Hu S, Hamer D. A functional polymorphism in the monoamine oxidase A gene promoter. *Hum Genet* 1998; 103(3): 273–279.
119. Scanlon SM, Williams DC, Schloss P. Membrane cholesterol modulates serotonin transporter activity. *Biochemistry* 2001; 40(35): 10507–10513.
120. Schalling D, Asberg M, Edman G, Oreland L. Markers for vulnerability to psychopathology: temperament traits associated with platelet MAO activity. *Acta Psychiatr Scand* 1987; 76(2): 172–182.
121. Schmidtke A, Bille-Brahe U, DeLeo D, Kerkhof A, Bjerke T, Crepet P et al. Attempted suicide in Europe: rates, trends and sociodemographic characteristics of suicide attempters during the period 1989–1992. Results of the WHO/EURO Multi-centre Study on Parasuicide. *Acta Psychiatr Scand* 1996; 93(5): 327–338.
122. Schmidtke A, Weinacker B, Fricke S. Epidemiologie von Suiziden und Suizidversuchen in Deutschland. *Suizidprophylaxe* 1998; Sonderheft: 37–49.
123. Schmitz G, Kaminski WE. ABC transporters and cholesterol metabolism. *Front Biosci* 2001; 6: D505–D514.
124. Schulsinger F. Psychopathy: heredity and environment. *Int J Ment Health* 1972; 1: 190–206.
125. Schulsinger F. A family study of suicide. In: Schov M, Stromgren G, editors. *Origin, Prevention and Treatment of Affective Disorders*. London, New York: Academic Press 1979: 277–287.
126. Sequeira A, Gwady FG, Ffrench-Mullen JM, Cagnetti L, Gingras Y, Casero RA, Jr et al. Implication of SSAT by gene expression and genetic variation in suicide and major depression. *Arch Gen Psychiatry* 2006; 63(1): 35–48.
127. Sequeira A, Kim C, Seguin M, Lesage A, Chawky N, Desautels A et al. Wolfram syndrome and suicide: Evidence for a role of WFS1 in suicidal and impulsive behavior. *Am J Med Genet B Neuropsychiatr Genet* 2003; 119B(1): 108–113.
128. Sequeira A, Mamdani F, Lalovic A, Anguelova M, Lesage A, Seguin M et al. Alpha 2A adrenergic receptor gene and suicide. *Psychiatry Res* 2004; 125(2): 87–93.
129. Serretti A, Calati R, Giegling I, Hartmann AM, Moller HJ, Rujescu D. Serotonin receptor HTR1A and HTR2C variants and personality traits in suicide attempters and controls. *J Psychiatr Res* 2008.
130. Serretti A, Mandelli L, Giegling I, Schneider B, Hartmann AM, Schnabel A et al. HTR2C and HTR1A gene variants in German and Italian suicide attempters and completers. *Am J Med Genet B Neuropsychiatr Genet* 2007; 144B(3): 291–299.
131. Sibille E, Arango V, Galfalvy HC, Pavlidis P, Erraji-Benchekroun L, Ellis SP et al. Gene expression profiling of depression and suicide in human prefrontal cortex. *Neuropsychopharmacology* 2004; 29(2): 351–361.
132. Skidgel RA, Erdos EG. The broad substrate specificity of human angiotensin I converting enzyme. *Clin Exp Hypertens A* 1987; 9(2–3): 243–259.
133. Stalenheim EG, von Knorring L, Oreland L. Platelet monoamine oxidase activity as a biological marker in a Swedish forensic psychiatric population. *Psychiatry Res* 1997; 69(2–3): 79–87.
134. Stanley M, Mann JJ. Increased serotonin-2 binding sites in frontal cortex of suicide victims. *Lancet* 1983; 1(8318): 214–216.
135. Statham DJ, Heath AC, Madden PA, Bucholz KK, Bierut L, Dinwiddie SH et al. Suicidal behaviour: an epidemiological and genetic study. *Psychol Med* 1998; 28(4): 839–855.
136. Stefulj J, Buttner A, Kubat M, Zill P, Balija M, Eisenmenger W et al. 5HT_{2C} receptor polymorphism in suicide victims. Association studies in German and Slavic populations. *Eur Arch Psychiatry Clin Neurosci* 2004; 254(4): 224–227.
137. Stefulj J, Buttner A, Skavic J, Zill P, Balija M, Eisenmenger W et al. Serotonin 1B (5HT_{1B}) receptor polymorphism (G861C) in suicide victims: association studies in German and Slavic population. *Am J Med Genet B Neuropsychiatr Genet* 2004; 127B(1): 48–50.
138. Strous RD, Bark N, Parsia SS, Volavka J, Lachman HM. Analysis of a functional catechol-O-methyltransferase gene polymorphism in schizophrenia: evidence for association with aggressive and antisocial behavior. *Psychiatry Res* 1997; 69(2–3): 71–77.
139. Strous RD, Nolan KA, Lapidus R, Diaz L, Saito T, Lachman HM. Aggressive behavior in schizophrenia is associated with the low enzyme activity COMT polymorphism: a replication study. *Am J Med Genet B Neuropsychiatr Genet* 2003; 120B(1): 29–34.
140. Swift RG, Sadler DB, Swift M. Psychiatric findings in Wolfram syndrome homozygotes. *Lancet* 1990; 336(8716): 667–669.
141. Thalmeier A, Dickmann M, Giegling I, Schneider B, Hartmann M, Maurer K et al. Gene expression profiling of post-mortem orbitofrontal cortex in violent suicide victims. *Int J Neuropsychopharmacol* 2008; 11(2): 217–228.
142. Traskman L, Asberg M, Bertilsson L, Sjostrand L. Monoamine metabolites in CSF and suicidal behavior. *Arch Gen Psychiatry* 1981; 38(6): 631–636.
143. Tripodianakis J, Markianos M, Sarantidis D, Spyropoulou G, Taktikou V, Bistolaki E. Platelet MAO activity in patients with dysthymic disorder. *Psychiatry Res* 1998; 78(3): 173–178.
144. Tsai SJ, Hong CJ, Hsu CC, Cheng CY, Liao WY, Song HL et al. Serotonin-2A receptor polymorphism

- (102T/C) in mood disorders. *Psychiatry Res* 1999; 87(2–3): 233–237.
145. Tsai SJ, Hong CJ, Yu YW, Chen TJ, Wang YC, Lin WK. Association study of serotonin 1B receptor (A-161T) genetic polymorphism and suicidal behaviors and response to fluoxetine in major depressive disorder. *Neuropsychobiology* 2004; 50(3): 235–238.
146. Tsai SJ, Wang YC, Hong CJ, Chiu HJ. Association study of oestrogen receptor alpha gene polymorphism and suicidal behaviours in major depressive disorder. *Psychiatr Genet* 2003; 13(1): 19–22.
147. Turecki G, Briere R, Dewar K, Antonetti T, Lesage AD, Seguin M et al. Prediction of level of serotonin 2A receptor binding by serotonin receptor 2A genetic variation in postmortem brain samples from subjects who did or did not commit suicide. *Am J Psychiatry* 1999; 156(9): 1456–1458.
148. Turecki G, Sequeira A, Gingras Y, Seguin M, Lesage A, Tousignant M et al. Suicide and serotonin: study of variation at seven serotonin receptor genes in suicide completers. *Am J Med Genet B Neuropsychiatr Genet* 2003; 118B(1): 36–40.
149. van den Oord EJ, Kuo PH, Hartmann AM, Webb BT, Möller HJ, Hettema JM, Giegling I, Bukszár J, Rujescu D. Genomewide association analysis followed by a replication study implicates a novel candidate gene for neuroticism. *Arch Gen Psychiatry* 2008; 65(9): 1062–71.
150. Verkes RJ, Fekkes D, Zwinderman AH, Hengeveld MW, van der Mast RC, Tuyl JP et al. Platelet serotonin and [3H]paroxetine binding correlate with recurrence of suicidal behavior. *Psychopharmacology (Berl)* 1997; 132(1): 89–94.
151. Verkes RJ, Pijl H, Meinders AE, van Kempen GM. Borderline personality, impulsiveness, and platelet monoamine measures in bulimia nervosa and recurrent suicidal behavior. *Biol Psychiatry* 1996; 40(3): 173–180.
152. Volavka J, Kennedy JL, Ni X, Czobor P, Nolan K, Sheitman B et al. COMT158 polymorphism and hostility. *Am J Med Genet B Neuropsychiatr Genet* 2004; 127B(1): 28–29.
153. Walther DJ, Peter JU, Bashammakh S, Hortnagl H, Voits M, Fink H et al. Synthesis of serotonin by a second tryptophan hydroxylase isoform. *Science* 2003; 299(5603): 76.
154. Wasserman D, Geijer T, Rozanov V, Wasserman J. Suicide attempt and basic mechanisms in neural conduction: relationships to the SCN8A and VAMP4 genes. *Am J Med Genet B Neuropsychiatr Genet* 2005; 133B(1): 116–119.
155. Wei J, Hemmings GP. Lack of evidence for association between the COMT locus and schizophrenia. *Psychiatr Genet* 1999; 9(4): 183–186.
156. Weinsilboum RM, Otterness DM, Szumlanski CL. Methylation pharmacogenetics: catechol O-methyltransferase, thiopurine methyltransferase, and histamine N-methyltransferase. *Annu Rev Pharmacol Toxicol* 1999; 39: 19–52.
157. Wender PH, Kety SS, Rosenthal D, Schulsinger F, Ortman J, Lunde I. Psychiatric disorders in the biological and adoptive families of adopted individuals with affective disorders. *Arch Gen Psychiatry* 1986; 43(10): 923–929.
158. Yanagi M, Shirakawa O, Kitamura N, Okamura K, Sakurai K, Nishiguchi N et al. Association of 14–3–3 epsilon gene haplotype with completed suicide in Japanese. *J Hum Genet* 2005; 50(4): 210–216.
159. Zalsman G, Braun M, Arendt M, Grunebaum MF, Sher L, Burke AK et al. A comparison of the medical lethality of suicide attempts in bipolar and major depressive disorders. *Bipolar Disord* 2006; 8(5 Pt 2): 558–565.
160. Zalsman G, Frisch A, Lewis R, Michaelovsky E, Hermesh H, Sher L et al. DRD4 receptor gene exon III polymorphism in inpatient suicidal adolescents. *J Neural Transm* 2004; 111(12): 1593–1603.
161. Zammit S, Jones G, Jones SJ, Norton N, Sanders RD, Milham C et al. Polymorphisms in the MAOA, MAOB, and COMT genes and aggressive behavior in schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 2004; 128B(1): 19–20.
162. Zhang HY, Ishigaki T, Tani K, Chen K, Shih JC, Miyasato K et al. Serotonin2A receptor gene polymorphism in mood disorders. *Biol Psychiatry* 1997; 41(7): 768–773.
163. Zhou Z, Roy A, Lipsky R, Kuchipudi K, Zhu G, Taubman J et al. Haplotype-based linkage of tryptophan hydroxylase 2 to suicide attempt, major depression, and cerebrospinal fluid 5-hydroxyindoleacetic acid in 4 populations. *Arch Gen Psychiatry* 2005; 62(10): 1109–1118.
164. Zill P, Buttner A, Eisenmenger W, Moller HJ, Bondy B, Ackenheil M. Single nucleotide polymorphism and haplotype analysis of a novel tryptophan hydroxylase isoform (TPH2) gene in suicide victims. *Biol Psychiatry* 2004; 56(8): 581–586.
165. Zill P, Preuss UW, Koller G, Bondy B, Soyka M. SNP- and haplotype analysis of the tryptophan hydroxylase 2 gene in alcohol-dependent patients and alcohol-related suicide. *Neuropsychopharmacology* 2007; 32(8): 1687–1694.
166. Zubenko GS, Maher BS, Hughes HB, III, Zubenko WN, Scott SJ, Marazita ML. Genome-wide linkage survey for genetic loci that affect the risk of suicide attempts in families with recurrent, early-onset, major depression. *Am J Med Genet B Neuropsychiatr Genet* 2004; 129B(1): 47–54.